

Identify Potential Treatment Lapses With MOUD Clarity™

Monitor potential lapses in treatment for opioid use disorder (MOUD) to improve patient care.

Patients who receive a MOUD after a non-fatal overdose significantly reduce their risk of an overdose death¹. However, given that treatments may be delivered over different periods of time and through multiple dispensations, providers may not always know when a patient discontinues an MOUD drug, which could increase the risk of a potential relapse or overdose. To improve outcomes for individuals receiving MOUD treatment during pivotal moments, providers need greater visibility into the continuity of MOUD treatment.

With MOUD Clarity™, providers can now seamlessly identify potential MOUD treatment lapses within existing workflows, delivering high-quality care with the right insights at the point of care.

What Is MOUD Clarity?

MOUD Clarity monitors dispensation data for drugs used to treat opioid use disorder to identify patients who may be experiencing a lapse in MOUD treatment. The enhancement automatically displays an indicator on the Patient Report and can be configured to send email communications and display dashboard alerts. MOUD Clarity empowers providers to know when to intervene in their patients' MOUD treatment to ensure continuity of care during pivotal care moments.

Did You Know?


drug overdose deaths in 20231 68.000




of those deaths had at least one potential opportunity for intervention²

more costly to care for patients 2.8-6.2x with behavioral health conditions including substance use disorder³

inih.gov/news-events/news-releases/methadone-buprenorphine-reduce-risk-death-after-opioid-overdose ²cdc.gov/overdose-prevention/data-research/facts-stats/sudors-dashboard-fatal-overdose-data.html iamanetwork.com/journals/jamanetworkopen/fullarticle/2799220





MOUD Clarity Empowers Providers With Knowledge During Pivotal Care Moments

MOUD Clarity allows providers to:



Identify patients who may be experiencing a lapse in MOUD treatment by checking if a patient has a dispensation gap greater than a specified threshold.



Receive notifications on potential gaps in MOUD treatment via emails sent to all providers and their delegates who have prescribed an MOUD drug to the patient in the last 45 days.



Easily view potential dispensation gap indicators displayed directly in PMP AWARxE® and PMP Gateway®.

To learn more about how your PDMP can improve care during pivotal care moments, see MOUD Clarity or contact us.